

---

# Taxonomy of *Basycladia* (Cladophorales, Chlorophyta) with Two New Combinations

David J. Garbary

Department of Biology, St. Francis Xavier University, Antigonish, Nova Scotia, Canada,  
B2G 2W5. dgarbary@gmail.com

---

**ABSTRACT.** The green algal genus *Basycladia* W. E. Hoffman & Tilden suggested for synonymy in *Cladophora* Kütz. by van den Hoek is reconfirmed as a genus based on the distinct basal system and the primarily epizoic habit associated with freshwater turtles. Previously assigned to *Cladophora* sect. *Basycladia* (W. E. Hoffmann & Tilden) C. Hoek, *C. kosteriae* C. Hoek (type: France) and *C. okamurae* (S. Ueda) C. Hoek (type: Japan) transfer to *Basycladia*, based on morphology and molecular evidence. The two new combinations *B. kosteriae* (C. Hoek) Garbary and *B. okamurae* (S. Ueda) Garbary bring the number of species in *Basycladia* to seven.

**Key words:** *Basycladia*, *Cladophora*, Cladophorales, turtle algae.

The cladophoroid genus *Basycladia* W. E. Hoffmann & Tilden (Hoffmann & Tilden, 1930) was initially described for two species of epizoic algae on the shells of freshwater turtles. Further species were added, all of which were either epizoic on turtles or occurred on a freshwater snail. Five species have now been assigned to the genus: *B. crassa* W. E. Hoffmann & Tilden (generitype), *B. chelonum* (Collins) W. E. Hoffmann & Tilden, *B. ramulosa* Ducker (Ducker, 1958), *B. sinensis* (N. L. Gardner) G. M. Sm. (Smith, 1950), and *B. vivipara* Normandin & Taft (Normandin & Taft, 1959). The genus *Basycladia* is widely distributed in freshwater habitats where turtles are present (e.g., Yoneda, 1952; Ducker, 1958; Prasad & Jain, 1973; Ernst & Norris, 1978; Garbary et al., 2007).

In his monograph of European *Cladophora* Kütz., van den Hoek (1963) considered *Basycladia* as a synonym of *Cladophora* and reduced *Basycladia* to sectional status within the genus. However, he incorrectly designated *B. chelonum* as the type, and none of the five existing species of *Basycladia* were formally transferred into *Cladophora*. A new species of *Cladophora*, *C. kosteriae* C. Hoek, was described, and *Chaetomorpha okamurae* S. Ueda was formally transferred to *Cladophora* as *C. okamurae* (S. Ueda) C. Hoek. Van den Hoek (1963) left us with several taxonomic and nomenclatural issues. First, the type of

*Cladophora* sect. *Basycladia* was not assigned to *Cladophora*. Furthermore, if *Basycladia* was distinct from *Cladophora* at generic rank, should *C. kosteriae* and *C. okamurae* continue to be considered as members of *Cladophora*, or should they be recognized as species of *Basycladia*? These issues can now be resolved.

Molecular evidence shows *Cladophora* to be extremely diverse, forming an ancient assemblage from which several green algal lineages are derived. Even within a limited clade (e.g., the *Cladophora albida* (Nees) Kütz.–*C. sericea* (Hudson) Kütz. series), there were six distinct ITS sequence types (Bakker et al., 1995). More recent molecular studies involving a diverse assemblage of Cladophorales and Siphonocladales showed that various siphonoclad lineages were associated with different clusters of *Cladophora* species (Leliaert et al., 2003, 2007). Thus, if numerous traditional genera of siphonoclad belonging to multiple families continue to be recognized, *Cladophora* can no longer be considered a natural genus.

A similar conclusion regarding the monophyly of *Cladophora* was inherent in the results of Yoshii et al. (2004) and Yoshii (2006), in which a wide range of species of Cladophorales was examined for pigment composition and 18S ribosomal DNA (rDNA) sequences. *Cladophora* species were sister to species in other genera of Cladophorales and Siphonocladales, e.g., *Chaetomorpha* Kütz., *Cladophoropsis* Børgesen, *Valonia* C. Agardh, and *Pithophora* Wittr. Hence, the cladophoroid morphology is plesiomorphic or possibly homoplasious within the assemblage, and *Cladophora* needs to be broken down into monophyletic units that can be recognized at generic level. Some of these units may already have been characterized in the various sections of *Cladophora* (van den Hoek, 1963).

One such segregate from *Cladophora* characterized by Yoshii et al. (2004) is *Basycladia*. While virtually all authors continue to recognize *Basycladia* (e.g., Ernst & Norris, 1978; Colt et al., 1995; John, 2003; Garbary et al., 2007), AlgaeBase follows van den Hoek (1963) and states that *Basycladia* is a synonym of *Cladophora* (Guiry & Guiry, 2007).



While the erect system of *Basycladia* with its branched, uniseriate filaments is similar to *Cladophora*, the poorly branched to unbranched, erect axes of *Basycladia* with very long initial cells in erect axes are characteristic. *Basycladia* species form a distinct basal layer of flattened, tightly packed cells, unlike the rhizoidal attachment of other *Cladophora* species. The association with turtles is diagnostic, even if not all species occur on turtles, and some species have been found on other substrata (Proctor, 1958; Normandin & Taft, 1959; van den Hoek, 1963). It is this heterotrichous basal layer that provides the basis for the association with turtles in that new erect filaments can be produced from a longer-lived basal system.

When van den Hoek (1963) reduced *Basycladia* to a section of *Cladophora* in his monograph of European *Cladophora*, none of the five species previously assigned to *Basycladia* were formally transferred. Furthermore, only two species were included in the section—the new species *C. kosteriae* and *C. okamuriae* (the latter transferred from *Chaetomorpha*; see Ueda, 1932). Neither species was found on turtles. Belusz and Reed (1969) later reported *C. kosteriae* from North America as epizoic algae on turtles. The association of *C. kosteriae* with *Basycladia* was later confirmed when Yoshii et al. (2004) used sequence data from type material of *C. kosteriae* to show a sister group relationship between *C. kosteriae* and two isolates of unidentified *Basycladia* from the United States in the UTEX culture collection (<<http://www.utex.org/>>).

Since *Basycladia* is a viable genus independent of *Cladophora*, and *C. kosteriae* is part of the *Basycladia* clade, *C. kosteriae* is hereby transferred to *Basycladia*. The morphology of *C. okamuriae* is also definitive for *Basycladia* and that species is also transferred.

1. ***Basycladia kosteriae*** (C. Hoek) Garbary, comb. nov. Basionym: *Cladophora kosteriae* C. Hoek, Revision of the European Species of *Cladophora* 1963: 37. TYPE: France. Paris: Jardin des Plantes, alpine garden, on stones in artificial freshwater stream, 25 Apr. 1961, C. van den Hoek n.61/9 (holotype, L not seen).
2. ***Basycladia okamuriae*** (S. Ueda) Garbary, comb. nov. Basionym: *Chaetomorpha okamuriae* S. Ueda, J. Imp. Fish. Inst., Tokyo 27: 23. 1932. *Cladophora okamuriae* (S. Ueda) C. Hoek, Revision of the European Species of *Cladophora* 1963: 39. TYPE: Japan. Tokyo: Shirahama (Bôshû), s.d., S. Ueda s.n. (holotype, Tokyo University of Marine Science and Technology not seen).

The genus *Basycladia* now comprises seven species: *B. crassa*, *B. chelonum*, *B. kosteriae*, *B. okamuriae*, *B. ramulosa*, *B. sinensis*, and *B. vivipara*. Guiry and Guiry (2007) list *B. sinensis* and *B. vivipara* as having provisional status. Neither of these species has been redescribed since their first descriptions. *Basycladia sinensis* is particularly ambiguous, as it was described from a turtle many months after it had been brought to California from China (Gardner, 1937). *Basycladia sinensis* is also morphologically similar to *B. okamuriae*. If these are conspecific, the latter has priority. *Basycladia vivipara* is known only from the freshwater snail *Viviparus malleatus* (Reeve). This apparent host specificity is analogous to that of the green alga *Sporocladopsis jackii* Garbary, C. J. Bird & K. Y. Kim (Garbary et al., 2005), which is known only from the marine snail *Ilyanassa obsoleta* (Say). Regardless, further studies are required on the status and relationships of *B. sinensis*, *B. okamuriae*, and *B. vivipara*.

*Acknowledgments.* This work was supported by grants from the Natural Sciences and Engineering Research Council of Canada to D.J.G.

#### Literature Cited

- Bakker, F. T., J. L. Olsen & W. T. Stam. 1995. Evolution of nuclear rDNA ITS sequences in the *Cladophora albida/sericea* clade (Chlorophyta). *J. Molec. Evol.* 40: 640–651.
- Belusz, L. C. & R. J. Reed. 1969. Some epizoophytes on six turtle species collected in Massachusetts and Michigan. *Amer. Midl. Naturalist* 81: 598–601.
- Colt, L. C., R. A. Saumure & S. Baskinger. 1995. First record of the algal genus *Basycladia* (Chlorophyta, Cladophorales) in Canada. *Canad. Field-Naturalist* 109: 454–455.
- Ducker, S. 1958. A new species of *Basycladia* on Australian freshwater turtles. *Hydrobiologia* 10: 157–174.
- Ernst, C. H. & J. N. Norris. 1978. Observations on the algal genus *Basycladia* and the Red-Bellied turtle *Chrysemys rubiventris*. *Estuaries* 1: 54–57.
- Garbary, D. J., C. J. Bird & K. Y. Kim. 2005. *Sporocladopsis jackii* sp. nov. (Chroolepidaceae, Chlorophyta): A new species from eastern Canada symbiotic with the mud snail, *Ilyanassa obsoleta* (Say) (Gastropoda). *Rhodora* 107: 52–68.
- , G. Bourque, T. B. Herman & J. A. McNeil. 2007. Epizoic algae from freshwater turtles in Nova Scotia. *J. Freshwater Ecol.* 22: 677–685.
- Gardner, N. L. 1937. A new species of *Chaetomorpha* from China. *Madroño* 4: 28–32.
- Guiry, M. D. & G. M. Guiry. 2007. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <<http://www.algaebase.org/>>, accessed 29 December 2007.
- Hirose, H. 1954. Studies on the morphology and behaviour of the reproductive cells of *Chaetomorpha okamurai* Ueda. *Cytologia* 19: 358–370.
- Hoffmann, W. E. & J. E. Tilden. 1930. *Basycladia*, a new genus of Cladophoraceae. *Bot. Gaz.* 89: 374–384.



- John, D. M. 2003. Filamentous and plantlike green algae. Pp. 311–352 in J. D. Wehr & R. G. Sheath (editors), *Freshwater Algae of North America: Ecology and Classification*. Academic Press, Amsterdam.
- Leliaert, F., F. Rousseau, B. De Reviere & E. Coppejans. 2003. Phylogeny of the Cladophorophyceae (Chlorophyta) inferred from partial LSU rRNA sequences: Is the recognition of a separate order Siphonocladales justified? *Eur. J. Phycol.* 38: 233–246.
- , O. De Clerck, H. Verbruggen, C. Boedeker & E. Coppejans. 2007. Molecular phylogeny of the Siphonocladales (Chlorophyta: Cladophorophyceae). *Molec. Phylogen. Evol.* 44: 1237–1256.
- Normandin, R. F. & C. E. Taft. 1959. A new species of *Basycladia* from the snail *Viviparus malleatus* Reeve. *Ohio J. Sci.* 59: 58–62.
- Prasad, B. N. & V. K. Jain. 1973. Observations on *Basycladia crassa* (Collins) Hoffmann and Tilden—A new addition to the Indian flora. *Curr. Sci.* 42: 363–364.
- Proctor, V. W. 1958. The growth of *Basycladia* on turtles. *Ecology* 39: 634–645.
- Smith, G. M. 1950. *Fresh-water Algae of the United States*, 2nd ed. McGraw-Hill, New York.
- Ueda, S. 1932. New freshwater species of *Chaetomorpha*. *J. Imp. Fish. Inst., Tokyo* 27: 23–24.
- Van den Hoek, C. 1963. *Revision of the European Species of Cladophora*. E. J. Brill, Leiden.
- Yoneda, Y. 1952. Observations on the algae growing on the pond tortoises, with species reference to *Basycladia crassa* Hoffmann & Tilden. *Publ. Seto Mar. Biol. Lab.* 2: 227–234.
- Yoshii, Y. 2006. Diversity and evolution of photosynthetic antenna systems in green plants. *Phycol. Res.* 54: 220–229.
- , T. Hanyuda, I. Wakana, K. Miyaji, S. Arai, K. Ueda & I. Inouye. 2004. Carotenoid compositions of *Cladophora* balls (*Aegagropila linnaei*) and some members of the Cladophorales (Ulvophyceae, Chlorophyta): Their taxonomic and evolutionary implication. *J. Phycol.* 40: 1170–1177.